

chemotaxis is inhibited as a consequence of increased macrophage spreading. Macrophage spreading is known to be one step of a series which leads to complete macrophage activation [28]. What this theory suggests is that TFF initiates an early phase in the sequence of macrophage activation and thereby immobilizes them. This results paradoxically in facilitation of tumor development.

Various degrees of macrophage activation have been defined. Thioglycollate-activated macrophages will spread rapidly but are not tumoricidal unless incubated with endotoxin [29]. However, macrophages stimulated with BCG, MAF, *Toxoplasma gondii*, or *Corynebacterium parvum* are cytotoxic to tumor cells including B16 [29-32]. Thioglycollate-induced peritoneal macrophages enhance B16 tumor development when injected s.c. with an inoculum of B16 cells [32]. While the exact level to which TFF activates the macrophage is yet to be determined, there is a precedent for partial activation of the macrophages.

It is not proved that the TFF and the macrophage spreading activity are the same molecule. However, there is a strong correlation between macrophage spreading and tumor facilitation. Both activities are systemic, and both are effective in nude mice. Tumor facilitation, inhibition of macrophage accumulation, and macrophage spreading are all induced by the same dose of material. In addition, the activities are present in both B16 culture supernatant and B16 tumor homogenate. The dose response curves for both tumor facilitation *in vivo*, and macrophage spreading *in vitro* show a supraoptimal range, in which the effects of these activities are reversed. Supraoptimal concentrations of TFF are not toxic to macrophages *in vitro* as measured by phagocytosis of sheep erythrocytes. Finally, the injection of viable B16 cells was previously [1] found to facilitate the development of tumors at distant sites. Macrophage spreading was also induced under these conditions. Thus, both TFF and macrophage spreading activity are produced by B16 cells *in vivo*. Although it is not proved that TFF and macrophage spreading activity are the same molecule, the correlation between these activities is strong.

REFERENCES

- Kalish RS, Brody NI: Increased incidence of tumor takes to B16 melanoma in C57B1/6J mice injected with B16 products. *J Invest Dermatol* 75:275-278, 1980
- Nakamura M, Ishida N, Kamo L: Immunosuppressive factors from mastocytoma cells cultured in serum-free medium. *J Natl Cancer Inst* 65:759-767, 1980
- Vandenbark AA, Burger DR, Finke P, Vetto RM: The effects of plasma from guinea pigs with tumor on PHA stimulated lymphocyte cultures. *Cell Immunol* 22:334-344, 1976
- Ting CC, Tsai SC, Rogers MJ: Host control of tumor growth. *Science* 197:571-573, 1977
- Ting CC, Rodrigues D, Ting RC: Suppression of T-cell mediated immunity by tumor cells: immunogenicity versus immunosuppression and preliminary characterization of the suppressive factors. *Int J Cancer* 24:644-655, 1979
- Kamo L, Patel C, Datey J, Friedman H: Immunosuppression induced *in vitro* by mastocytoma tumor cells and cell-free extracts. *J Immunol* 114:1749-1756, 1975
- Glasgow AH, Nimberg RB, Menzoian JO, Saporoschetz I, Cooperband SR, Schmid K, Mannick JA: Association of anergy with an immunosuppressive peptide fraction in the serum of patients with cancer. *N Engl J Med* 291:1263-1267, 1974
- Nowothny AR, Butler C, Grohman J, Keebler C: Dual effects of tumor antigens: induction of tumor resistant or tumor growth enhancement. *Ann NY Acad Sci* 276:106-123, 1976
- Gresser I, Vignaux F, Maury C, Lindahl PI: Factor(s) from Ehrlich ascites cells responsible for delayed rejection of skin allografts in mice and its assay on lymphocytes *in vitro*. *Proc Soc Exp Biol Med* 149:83-88, 1975
- North RJ, Kirstein DP, Tuttle RL: Subversion of host defense mechanisms by murine tumors. I. A circulating factor that suppresses macrophage-mediated resistance to infection. *J Exp Med* 143:559-573, 1976
- Jessup JM, Cohen MH, Tomaszewski MM, Felix EL: Effects of murine tumors on delayed hypersensitivity to dinitrochlorobenzene. I. Description of anergy caused by transplanted tumors. *J Natl Cancer Inst* 57:1077-1082, 1976
- Nelson M, Nelson DS: Macrophages and resistance to tumors. I. Inhibition of delayed-type hypersensitivity reactions by tumor cells and by soluble products affecting macrophages. *Immunology* 34:277-289, 1978
- Nelson M, Nelson DS: Macrophages and resistance to tumors. IV. Influence of age on susceptibility to anti-inflammatory and anti-macrophage effects of tumor cell products. *J Natl Cancer Inst* 65:781-789, 1980
- Normann SJ, Sorkin E: Inhibition of macrophage chemotaxis by neoplastic and other rapidly proliferating cells *in vitro*. *Cancer Res* 37:705-711, 1977
- Snyderman R, Pike MC: An inhibitor of chemotaxis produced by neoplasms. *Science* 192:370-372, 1976
- Cianciolo GJ, Herberman RB, Snyderman R: Depression of murine macrophage accumulation by low-molecular-weight factors derived from spontaneous mammary carcinomas. *J Natl Cancer Inst* 65:829-834, 1980
- Snyderman R, Pike MC, Blaylock BL, Weinstein P: Effects of neoplasms on inflammation: depression of macrophage accumulation after tumor implantation. *J Immunol* 116:585-589, 1976
- James K: The influence of tumor cell products on macrophage function *in vitro* and *in vivo* (review), Proceedings of the EURES Symposium on the Macrophage and Cancer, Edinburgh, 1977. Edited by K James, B McBride, A Stuart. pp 225-246
- Pike MC, Snyderman R: Depression of macrophage function by a factor produced by neoplasms: a mechanism for abrogation of immunosurveillance. *J Immunol* 117:1243-1249, 1976
- Cheung HT, Cantarow WD, Sundharadas G: Characteristics of a low-molecular-weight factor extracted from mouse tumors that affects *in vitro* properties of macrophages. *Int J Cancer* 23:344-352, 1979
- Lowry OH, Rosenbrough NO, Farr L, Randall RJ: Protein measurement with the Folin phenol reagent. *J Biol Chem* 193:265-275, 1951
- Bianco C, Eden A, Cohn ZA: The induction of macrophage spreading: role of coagulation factors and the complement system. *J Exp Med* 144:1531-1544, 1976
- Davis BJ: Disc electrophoresis. II. Method and application to human serum proteins. *Ann NY Acad Sci* 121:404-427, 1964
- Prenn LM, Outzen HC: Primary tumor immunity in nude mice. *Int J Cancer* 19:688-691, 1977
- Warner NL, Woodruff MFA, Burton RC: Inhibition of the growth of lymphoid tumors in syngeneic athymic (nude) mice. *Int J Cancer* 20:146-155, 1977
- Meltzer MD: Tumoricidal responses *in vitro* of peritoneal macrophages from conventionally housed and germ-free nude mice (short communication). *Cell Immunol* 22:176-181, 1976
- Herberman RB, Nunn ME, Lavin DH: Natural cytotoxic reactivity of mouse lymphoid cells against syngeneic and allogeneic tumors. I. Distribution of reactivity and specificity. *Int J Cancer* 16:216-219, 1975
- Blanden RV: Modification of macrophage function. *J Reticuloendothel Soc* 5:179-202, 1968
- Hibbs JB, Taintor RR, Chapman HA, Weinberg JB: Macrophage tumor killing: influence of the local environment. *Science* 197:279-282, 1977
- Leonard EJ, Ruco LP, Meltzer MD: Characterization of macrophage activation factor, a lymphokine that causes macrophages to become cytotoxic for tumor cells. *Cell Immunol* 41:347-357, 1978
- Moore K, McBride WH: The activation state of macrophage subpopulations from a murine fibrosarcoma. *Int J Cancer* 26:609-615, 1980
- Gabizon A, Lelbovich SJ, Goldman R: Contrasting effects of activated and non-activated macrophages from tumor bearing mice on tumor growth *in vivo*. *J Natl Cancer Inst* 65:913-920, 1980